

# What is Citizen Science?

Also called Citizen Participatory Research





# It is a partnership in which public participants are involved in a scientific endeavor.

Partnerships can occur between:

- a university and a watershed group (e-Bird: Cornell and bird counts)
- a watershed group and its community (SYRCL and Nat. park closure)
- a scientific institute and its community (SSI and breast cancer study)
- a government agency and a watershed group (Safe to Swim).

# Citizen Monitoring - Who is involved?





There are over 200 watershed groups in California, so many opportunities for partnerships



**Examples of watershed group projects** 

**Monitoring**- Stormwater, First Flush, BMP Assessment, Benthic Macroinvertebrate and Algae Bioassessments, TMDLs, Stormwater, 303(d) listings, Secchi Dip-In



**Community Health** - mining contaminants, *E. coli*, Safe to Eat






**Education** —public & charter schools STEM, Safe to Swim Fish contamination, trash



**Restoration -** native vegetation, salmon, Brownfields mining








 Started in 1996 as Friends of Deer Creek by a group of concerned local citizens and property owners, when Nevada City was replacing the historic Pine Street Bridge.

 Focus on scientific investigation and methods, to find solutions to Deer Creek's problems.



# **Deer Creek Watershed**



# **Community-based Participatory Research**

### Sierra Streams Institute is working with local citizens to improve:

- environmental health of ecosystems
- human health of community members
- science/public health education

Citizens participate in all levels of work.



### Some Current Projects:

- Restoration of native vegetation and endangered salmon habitate
- Health impacts/remediation of mining contaminants
- Health impacts/source id of bacterial contamination in swimming holes
- Study of correlation of Nisenan childhood stress with current health

# Providence Mine – Then...




Photo 50. Providence Mine, Nevada City District. This 1893 view of the mine, in Nevada County, looks southeast. The Champion mine is at left, Deer Creek in the foreground.



# EPA Brownfields Assessment/Cleanup




Cleanup includes traditional dig and haul and innovative methods such as:

Phytoremediation and addition of fish bone meal.



# Why monitor? *E. coli* contamination






# **Pioneer Park Restoration**



- Could not remove cement due to Arsenic contamination
- Removed invasive plant species, replanted with natives
- Added boulders and shifted bridge to change hydrology, reduce sedimentation
- Successful at getting City of Nevada City to follow BMP's

# Pioneer Park Restoration- BMI



Axis 1

# **Indicator Species Analysis**

(Dufrene and Legendre 1997)

## **Group 1 (Before Restoration)**

Trichoptera, Polycentropodidae: Indicator Value= 32.1, p= 0.21 Filterer-Collector, Tolerance Value= 6

# **Group 2 (After Restoration)**

Megaloptera, Corydalidae: Indicator Value= 37.5, p= 0.2 Predator, Tolerance Value= 0

Plecoptera, Pteronarcyidae: Indicator Value= 40, p= 0.17 Shredder, Tolerance Value = 0



# Summary

- Restoration work is improving conditions for BMI communities
- BMI are good indicators of ecosystem health
- Results show volunteers & community how successful their work has been




# Lake Wildwood Wastewater Treatment Plant Upgrade

- Government mandate in June 2007
- Upgrade to fully denitrify wastewater, produce more consistent, contained flows
- Reduced NO<sub>3</sub> from  $\mu$ = 1.085 mg/L to 0.67 mg/L (SE ± 0.18, z= -440.5, p= 0.03)
- Reduced water temperature from  $\mu$ = 18.5°C to 15.9°C (SE ± 1.24, t= 2.16, p= 0.04)



# LWW Upgrade- BMI Communities



p = 0.057

A = 0.006



Indicator Species Analysis (Dufrene and Legendre 1997)

**Group 1 (Before Upgrade)** 

None!

### **Group 2 (After Upgrade)**

Diptera, Tipulidae: Indicator Value= 30.2, p= 0.05 Shredder Tolerance Value= 3



# **Education Program**





- Scientist led
- Biological Field Sampling and Monitoring
- Laboratory Experiences
- Science Career Panels
- Student written Field Guides
- Native plant Restoration



# The 2<sup>nd</sup> edition of THE BUG BOOK is now on sale. Order by calling 530 265-6090

#### Order Summary

Match your sample to the images and descriptions below.

### COLEOPTERA



- Segmented legs Leathery plates on entire body
- Hard, oval shaped
- Maxillary palps and antenna extending from head

### COLEOPTERA Adults



DIPTERA

Soft, fleshy body with

### EPHEMEROPTERA



- 1 set of wingpads Segmented legs with 1
- claw on each
- 2 or 3 tails

### Chapter 4



· Head prominent with large eyes

**ODONATA** 

Chapter 7

- Wingpads
- Segmented legs

#### HEMIPTERA Adults Chapter 4



Ends of wings membranous



- Prominent head Short, stubby thoracic legs fused to body
- Prolegs on abdomen

### Chapter 6



- · Body slightly flattened
- with big mouthparts Segmented legs, various tails and append-

NON-INSECTS

Chapter 10



- 2 pairs of wingpads Large head with segmented antenna, large eyes
- · Segmented legs with 2 claws
- Abdomen ends in short spikes or flat, thin gills

#### PLECOPTERA Chapter 8



- Long antennae 2 long, thin tails
- · 2 pairs of wingpads Segmented legs with



May be in a case Segmented legs

xviii

2 hooks on prolegs on end of abdomen







With 4 or more pairs of jointed legs

Friends of Deer Creek Nevada City, California www.friendsofdeercreek.org

Order: Suborder: Family: # Genera/Species: Common Name:

Odonata Zygoptera COENAGRIONIDAE

NA: 14 gen, 116 spp; CA: 8 gen Narrow-winged or Pond Damselflies

#### Taxonomic Characteristics:

- Body length 13–25 mm
- All segments of antennae about the same length (b)
- Labium (lower lip) stout and somewhat trapezoidal without medial notch (d)
- 3 inflated, leaf-like gills similar in length, ending in a point, and with small, highly branched veins joining main vein diagonally

### **Biological Information:**

Tolerance Value: 9

Functional Feeding Group: p

Coenagrionidae are the largest family of damselflies in North America. Most prefer the slow, quiet water of ponds and stream edges where the larvae climb on vegetation or sprawl on soft bottom sediment, while a few species favor pool or riffle areas in streams or



a. Argia, dorsal view







a-b. McCormick; c. Elder; d-e. McCormick


c. Leaf-like gills

Taxa Tips! Easily identified by ruling out other families. Calopterygidae have large basal antennal segments that are longer than all the others combined, and Lestidae have an extremely long, narrow labium.

7-5

Notes:

a. McCormick; b. CDFG ABL; c. McCormick; d. Cutter, L.; e-f. Elder; g. McCormick; h. Elder; i. McCormick; j. Elder; k-l. McCormick; m-n. Elder



www.sierrastreams.org



Linking water, science, and people.